

# Carbo K (0-0-21)

Potassium containing liquid plant nutrient for foliar feeding to maintain or correct potassium levels.

## GUARANTEED ANALYSIS:

Soluble Potash (K<sub>2</sub>O) .....21%

Derived from: Potassium Acetate

## NOTE:

This product should be used as part of a complete fertilizer program. This fertilizer should be used only as recommended. Product may be harmful if misused.

Information regarding the contents and levels of metals in this product is available on the internet at:  
<http://www.aapfco.org/metals.htm>

## EMERGENCY TELEPHONE:

Chemtrec U.S.-Canada: 800-424-9300

Chemtrec International: 703-527-3887

☐ **2.5 gallons (9.46 ℓ)**  
**Net Weight: 26.50 lbs (12.01 kg)**

☐ **2 X 2.5 gallons (2 x 9.46 ℓ)**  
**Net Wt. 53 lbs. (24.02 kg)**

☐ **55 gallons (208.19 ℓ)**  
**Net Wt. 583 lbs (264.4 kg)**

☐ **275 gallons (1040.99 ℓ)**  
**Net Wt. 2915 lbs (1322 kg)**

**Density 10.60 lbs/gal at 68°F**

CONDITIONS OF SALE AND WARRANTY: HydroGro (Seller) warrants that this product conforms to the chemical description on the label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS EXPRESSED, IMPLIED, OR STATUTORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. No one is authorized to make any other warranty, guarantee, or directions concerning this product, and no such warranties, guarantees or directions shall be valid or binding upon Seller. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or an accompanying written recommendation, are beyond the control of Seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. In no case shall Seller be liable for consequential, special, indirect, or incidental damages or losses resulting from the handling or use of this product. The foregoing is a condition of sale by Seller and is accepted as such by the buyer. Buyer further agrees in the event of damages arising from the use of this product to accept a replacement of the product or a refund of the purchase price of the product, at buyer's option, as full discharge of Seller's liability.



## WARNING

### HAZARD STATEMENTS:

Harmful if swallowed.  
Causes skin irritation.  
Causes serious eye irritation.  
May cause respiratory irritation.

### PRECAUTIONARY STATEMENTS

#### PREVENTION:

Avoid breathing dust/ fume/gas/mist/vapors/spray.  
Wash skin thoroughly after handling.  
Do not eat, drink, or smoke when using this product.  
Use only outdoors or in a well-ventilated area.  
Wear protective gloves/protective clothing/  
eye protection/ face protection.

#### RESPONSE:

IF SWALLOWED: Call a poison center or doctor/  
physician if you feel unwell. Rinse mouth.  
IF ON SKIN: wash with plenty of soap and water.  
IF INHAILED: Move victim to fresh air and keep  
at rest in a comfortable position for breathing.  
IF IN EYES: Rinse cautiously with water for several  
minutes. Remove contact lenses, if present an easy to  
do. Continue rinsing.  
If skin irritation occurs: Get medical advice / attention  
Take off contaminated clothing.

## DIRECTIONS FOR USE :

Shake well before use.

### TANK MIXING:

Read ALL labels carefully and adhere strictly to the instructions for use and advice regarding whether or not product(s) should be co-applied. Many variables can influence the performance of co-applied products and therefore co-application is entirely at the risk of the end-user. It is strongly recommended that a limited application is made initially when using unfamiliar product mixes. Before co-application of products you, or your advisor, must do a jar test.

### CAUTION:

READ THE LABEL BEFORE USING.  
KEEP OUT OF REACH OF CHILDREN AND PETS.

### STORAGE:

Store locked up

### DISPOSAL:

Dispose of waste materials and container by following the waste disposal requirements of your country, state, or local authorities



# Carbo K (0-0-21)

## DIRECTIONS FOR USE:

*Use only as Directed.*

*Carbo K* should only be used where there is a need. Do not exceed appropriate application rates. Avoid application under climate extremes.

*Carbo K* could lower the pH of the spray water, therefore the use of a buffer is recommended when using in combination with other chemicals.

*Carbo K* should be applied in the early morning or late afternoon. Do not apply at temperatures above 85°F. Do not apply to plants that are undergoing a period of moisture or heat stress.

CROP	USAGE RATE	REMARKS
<b>GRAPES</b> Table, wine, raisins	1 pt - 1.5 gal	Apply 1-3 times, starting at pea size. Repeat every 10-14 days. Apply 80-100 gallons per acre. <b>Table Grapes:</b> Do not apply within 4 weeks of harvest.
<b>BERRIES</b> Blueberries, Raspberries, Strawberries	1 pt - 1 gal	Apply during fruit fill and ripening at 10-14 day intervals as required. Based on a typical water volume of 25-150 gallons per acre.
<b>CEREALS</b> Wheat, barley, oats	1 qt - 1.5 gal	Apply during the reproductive stage of the plants.
<b>CANOLA AND SOYBEANS</b>	1 qt - 1.5 gal	Apply during the reproductive stage of the plants.
<b>STONE AND POME FRUIT</b> Plums, peaches, olives, cherries, apricots	1 pt - 2 gal	Apply 2-3 applications after petal fall. <b>Fruit color and size:</b> Apply the first application 4 weeks prior to harvest and the second application 2 weeks later. Apply another 1-2 applications post-harvest.
<b>SUBTROPICAL FRUIT</b> Avocados, citrus, coffee	2 qts - 1.5 gal	Apply 1-2 applications between bud break and the onset of flowering (two-thirds of growth flush completed). Apply 1-2 applications after petal fall. <b>Fruit color and size:</b> Apply the first application 4 weeks prior to harvest and the second application 2 weeks later. Apply another 1-2 applications post-harvest.
<b>NUTS</b> Pecans, walnuts, almonds, cashews, pistachio, macadamias	1 pt - 2 gal	Apply the first application in the spring and then on all new growth later in the season or when deficiencies occur. 2-3 applications are recommended. Apply in 100-200 gallons of water per acre.
<b>VEGETABLES</b>	1 pt - 1 gal	Apply 2-3 weeks after germination and repeat 3-4 weeks later if necessary. Apply in 25-125 gallons of water per acre.

