Derived from: Zinc gluconate

complete fertilizer program. This

levels of metals in this product is

http://www.aapfco.org/metals.html.

available on the internet at:

EMERGENCY TELEPHONE:

2.5 gallons (9.46 ℓ)

Net Wt. 29.2 lbs (13.2 kg)

2 X 2.5 gallons (2 x 9.46 l)

Net Wt. 58.4 lbs (26.5 kg)

Net Wt. 350.4 lbs (158.9 kg)

30 gallons (113.56 ℓ)

275 gallons (1040.99 ℓ) Net Wt. 3,212 lbs (1,457 kg)

fertilizer should be used only as

This product should be used as part of a

recommended. Product may be harmful

Information regarding the contents and

Chemtrec U.S.-Canada: 800-424-9300

NOTE:

if misused.

A liquid fertilizer for use as a foliar feed to maintain or correct zinc levels in plants.



DANGER







HAZARD STATEMENTS:

Causes serious eye damage.
May damage fertility or unborn child.
Very toxic to aquatic life.
Very toxic to aquatic life with long lasting effects.

PRECAUTIONARY STATEMENTS

Wear protective gloves/protective clothing/eye protection/face protection.

IF IN EYES: Immediately rinse with water for several minutes. Remove contact lenses, if present and easy to do. Continue rinsing. Get medical help. If exposed or concerned, get medical advice.

PREVENTION:

Wash hands and affected area thoroughly after handling.
Do not touch eyes.
Avoid release to the environment.

RESPONSE:

The symptoms resulting from direct exposure to the product could appear a while after exposure. If there is persistent discomfort, seek medical attention. Immediately remove contaminated clothing and remove the affected person from the contamination area.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED READ THE LABEL BEFORE USING.

KEEP OUT OF REACH OF CHILDREN AND PETS.

STORAGE:

Store in a well ventilated place. Keep container tightly closed.
Avoid direct sunlight.
Store locked up.

DISPOSAL:

Dispose of waste materials and container by following the waste disposal requirements of your country, state, or local authorities.

Density 11.68 lbs/gal at 68°F





A liquid fertilizer for use as a foliar feed to maintain or correct zinc levels in plants.

DIRECTIONS FOR USE: USE ONLY AS DIRECTED.

- For optimum results, GLUCO-ZINC should be used in a program and in combination with other balanced fertilizers and foliar feeds.
- GLUCO-ZINC is compatible with most agricultural chemicals. It is advisable to do a miscibility (jar) test prior to mixing.
- Do NOT mix GLUCO-ZINC with highly alkaline material such as Lime Sulfur and Bordeaux mixture or with any phosphate containing fertilizers.
- GLUCO-ZINC should preferably be applied in the early morning or late afternoon. Do NOT apply to plants that are undergoing a period of
 moisture or heat stress.
- Consult with your chemical distributor should crop specific programs or any other information be required about the use of GLUCO-ZINC.
- Use lower rates as a maintenance spray and higher rates to correct deficiencies.

CROP	MAX RATE/A	REMARKS
CITRUS		Apply as a full cover spray in spring to all new growth (2-3 applications). Do not spray directly before or during harvest. Apply in 200 gal water per acre.
CORN	28 oz	Make first application early in the season at the first signs of a deficiency and repeat if required. Apply in 20 gal water per acre.
ONIONS	2 qt	Apply at the first signs of a deficiency. Repeat application 2 weeks later if required. Apply in 50 gal water per acre.
POTATOES	28 oz	Apply at the first signs of a deficiency and repeat 3-4 weeks later if necessary. Apply in 50 gal water per acre.
WHEAT/BARLEY	28 oz	Apply early in the season at the first signs of a deficiency and repeat the application if required. Apply in 20 gal water per acre.
APPLES/PEARS	28 oz	2 applications per season. 1 early in the season and again after harvest in 50-200 gal water per acre.
BEANS, GROUNDNUTS, PEAS, SOYBEANS	28 oz	1-2 applications early in the growing season. Apply in 20 gal water per acre.
OTHER CROPS	14 oz	Apply as a full cover spray and repeat every 14 days as needed in 50 gal water per acre.

CONDITIONS OF SALE AND WARRANTY: HydroGro (Seller) warrants that this product conforms to the chemical description on the label. THIS WARRANTY IS EXPRESSLY IN LIEU OF ALL OTHER WARRANTIES AND REPRESENTATIONS EXPRESSED, IMPLIED, OR STATUATORY, INCLUDING WARRANTIES OF MERCHANTABILITY AND FITNESS FOR A PARTICULAR USE. No one is authorized to make any other warranty, guarantee, or directions concerning this product, and no such warranties, guarantees or directions shall be valid or binding upon Seller. Timing, rate and method of application, weather and crop conditions, mixtures with chemicals not specifically recommended on this label or an accompanying written recommendation, are beyond the control of Seller. Buyer assumes all risks of use, storage and handling of this material not in strict accordance with directions given herewith. In no case shall Seller be liable for consequential, special, indirect, or incidental damages or losses resulting from the handling or use of this product. The foregoing is a condition of sale by Seller and is accepted as such by the buyer. Buyer further agrees in the event of damages arising from the use of this product to accept a replacement of the product or a refund of the purchase price of the product, at buyer's option, as full discharge of Seller's liability.

